arXiv:1302.6187v1 [nlin.CD] 25 Feb 2013

The relevance of Brody level spacing distribution in dynamically localized chaotic eigenstates

Benjamin Batistić*, Thanos Manos*,† and Marko Robnik*

*CAMTP - Center for Applied Mathematics and Theoretical Physics, University of Maribor, Krekova 2, SI-2000 Maribor, Slovenia, European Union

† School of Applied Sciences, University of Nova Gorica, Vipavska 11C, SI-5270 Ajdovščina, Slovenia, European Union

E-mail: Benjamin.Batistic@gmail.com, Thanos.Manos@uni-mb.si, Robnik@uni-mb.si

Abstract. We study dynamically localized chaotic eigenstates in the finite dimensional quantum kicked rotator as a paradigm of Floquet systems and in a billiard system of the mixed-type (Robnik 1983) as a paradigm of time-independent Hamilton systems. In the first case we study the spectrum of quasienergies, in the second one the energy spectrum. In the kicked rotator we work in the entirely chaotic regime at $K = 7$, whilst in the billiard we use the Poincaré Husimi functions (on the Poincaré Birkhoff surface of section) to separate the regular and chaotic eigenstates, and then perform the analysis of 587654 high-lying chaotic eigenstates (starting at about 1.000.000 above the ground state). In both cases we show that the Brody distribution excellently describes the level spacing distribution, with an unprecedented accuracy and statistical significance. The Berry-Robnik picture of separating the regular and chaotic levels in the case of the billiard is also confirmed.

PACS numbers: 01.55.+b,02.50.Cw,02.60.Cb,05.45.Pq, 05.45.Mt, 47.52.+j

Submitted to: *J. Phys. A: Math. Gen.*

1. Introduction

Quantum chaos is still an important and lively field of research, especially as it is moving into more and more difficult experimental and numerical investigations. The study of a deep semiclassical regime requires great experimental and numerical algorithmical and technical efforts to reach sufficiently many accurate high-lying eigenstates, and still uncovers surprising phenomena. In (general) generic time-independent (autonomous) Hamilton systems [1, 2, 3] in the strict semiclassical limit we can conceptually separate regular and chaotic eigenstates, the classification being based on the Wigner functions [4] of the eigenstates, or other quantal phase space functions like the Husimi functions [5] (Gaussian smoothed Wigner functions). This picture goes back to the pioneering ideas by Percival in 1973 [6] and to the first theoretical work by Berry and Robnik in 1984 [7], and has been perhaps most clearly manifested in the work of Prosen and

Robnik [8, 9, 10], see also [11, 12], and in the recent work of Batistić and Robnik [13], [14].

One of the main discoveries in quantum chaos is the finding that in classically fully chaotic, ergodic, autonomous Hamilton systems with the purely discrete spectrum the fluctuations of the energy spectrum around its mean behaviour obey the statistical laws described by the Gaussian Random Matrix Theory (RMT) [1, 2, 15, 16], provided that we are in the sufficiently deep semiclassical limit. The latter condition means that all relevant classical transport times, like the typical ergodic time, or diffusion time, are smaller than the so-called Heisenberg time, or break time, given by $t_H = 2\pi\hbar/\Delta E$, where $h = 2\pi\hbar$ is the Planck constant and ΔE is the mean energy level spacing, such that the mean energy level density is $\rho(E) = 1/\Delta E$. This statement is known as the Bohigas - Giannoni - Schmit (BGS) conjecture and goes back to their pioneering paper in 1984 [17], although some preliminary ideas were published in [18]. Since $\Delta E \propto \hbar^d$, where d is the number of degrees of freedom (= the dimension of the configuration space), we see that for sufficiently small $\hbar$ the stated condition will always be satisfied. Alternatively, fixing the $\hbar$, we can go to high energies such that the classical transport times become smaller than t_H. The role of the antiunitary symmetries that classify the statistics in terms of GOE, GUE or GSE (ensembles of RMT) has been explained in [19], see also [20] and [1, 2, 3, 15]. The theoretical foundation for the BGS conjecture has been initiated first by Berry [21], using the Gutzwiller periodic orbit theory (trace formula) [22] (for an excellent exposition see [1]) and later further developed by Richter and Sieber [23], arriving finally in the almost-final proof proposed by the group of F. Haake [24, 25, 26, 27].

However, quite generally, if the semiclassical condition is not satisfied, such that t_H is no longer larger than the relevant classical transport time, like e.g. the diffusion time in fully chaotic but slowly ergodic systems, we find the so-called **dynamical localization**, or **Chirikov localization**. Dynamical localization was first discovered in time dependent systems [28]. It was intensely studied since then in particular by Chirikov, Casati, Izrailev, Shepelyanski and Guarneri, in the case of the kicked rotator as reviewed in [29]. See also the references [30]-[33], and the most recent work [34]. For a general overview of the time periodic (Floquet) systems see also [1, 2]. It has been observed that in parallel with the localization of the eigenstates one finds the fractional power law level repulsion (of the quasienergies) even in fully chaotic regime (of the finite dimensional kicked rotator), and it is believed that this picture applies also to time independent (autonomous) Hamilton systems and their eigenstates [34]. (See the excellent review of localization in time independent billiards by Prosen in [35].) Indeed, this has been analyzed indirectly with unprecedented precision and statistical significance recently by Batistić and Robnik [13] in case of mixed-type systems. The present work concerns the theoretical separation of the regular and chaotic eigenstates and the analysis of the chaotic eigenstates and the corresponding energy subspectrum. The details will be published in a separate paper [14]. An early attempt of separation of eigenstates in the billiard system has been published in [36], using a different approach at much lower energies and with much smaller statistical significance.

In case that the system is classically integrable, Poisson statistics applies, as is well known and goes back to the work by Berry and Tabor in 1977 (see [1, 2, 3] and the references therein, and for the recent advances [37]).

In the mixed-type regime, where classical regular regions coexist in the classical phase space with the chaotic regions, exhibiting a typical KAM-scenario which is the generic situation, the Principle of Uniform Semiclassical Condensation (of the Wigner functions of the eigenstates; PUSC) applies, based on the ideas by Berry [38], and further extended by Robnik [3]. The Wigner functions uniformly condense on the classical invariant component. Then, if the stated semiclassical condition is satisfied, the chaotic eigenstates are uniformly extended chaotic, and consequently the Berry-Robnik statistics [7, 10] is observed - see also [3]. If the semiclassical condition stated above requiring that t_H is larger than all classical transport times is not satisfied, the chaotic eigenstates will not be extended but localized, dynamically localized, instead and the Berry-Robnik statistics must be generalized as explained in [8, 9, 3, 39, 13, 14], in order to capture the localization effects. Namely, instead of GOE for the chaotic eigenstates we must allow for the Brody distribution [40, 41] and this is the main message of the present work. The Brody level spacing distribution is defined as

$$P_B(S) = C_1 S^{\beta} \exp\left(-C_2 S^{\beta+1}\right), \quad W_B(S) = 1 - \exp\left(-C_2 S^{\beta+1}\right), \tag{1}$$

where the two parameters C_1 and C_2 are determined by the two normalizations $<1>=<S>=1$, and are given by

$$C_1 = (\beta+1)C_2, \quad C_2 = \left(\Gamma\left(\frac{\beta+2}{\beta+1}\right)\right)^{\beta+1}, \tag{2}$$

with $\Gamma(x)$ being the Gamma function. By $W_B(S)$ we denote the cumulative Brody level spacing distribution, $W(S) = \int_0^S P(x)\, dx$. The Brody parameter β is in the interval $[0,1]$, where $\beta = 0$ yields the Poisson distribution in case of the strongest localization, and $\beta = 1$ gives the Wigner surmise (2D GOE, as an excellent approximation of the infinite dimensional GOE), which describes the extended chaotic eigenstates. It turns out that the Brody distribution fits the empirical data much better than e.g. the distribution function proposed by F. Izrailev (see [30, 33, 29] and the references therein), defined as

$$P_I(S) = A\left(\frac{1}{2}\pi S\right)^{\beta} \exp\left[-\frac{1}{16}\beta\pi^2 S^2 - \left(B - \frac{1}{4}\pi\beta\right)S\right], \tag{3}$$

where the constants A and B are determined by the normalizations $<1>=<S>=1$.

If in a mixed-type system the couplings between the regular eigenstates and chaotic eigenstates become important, at low energies, due to the dynamical tunneling, we can use the ensembles of random matrices that capture these effects [42, 43, 44, 13]. As the tunneling strengths typically decrease exponentially with the inverse effective Planck constant, they rapidly disappear with increasing energy, or by decreasing the value of the Planck constant. In this work we shall deal only with high-lying eigenstates, and therefore we can neglect the effects of tunneling.

In this paper, in section 2, we introduce a criterion for classifying eigenstates as regular and chaotic in the billiard system [45, 46], and use it to separate the regular and chaotic eigenstates as well as the corresponding eigenenergies, and moreover, we show that the regular levels obey the Poisson statistics, whilst the chaotic dynamically localized eigenenergies obey exceedingly well the Brody distribution (1). This we do with great accuracy and statistical significance. In section 3 we consider a finite dimensional kicked rotator model [30, 33, 29], which is in fact also the quantized standard or Chirikov map, and show again that the level spacings between the quasienergies of this Floquet system obey the Brody distribution better than Izrailev's one. It is well known that Brody distribution so far has no theoretical foundation, but our empirical results show that we have to consider it seriously in dynamically localized chaotic eigenstates, thereby being motivated for seeking its physical foundation. The details of this study will be published in separate papers, namely in [34] for the kicked rotator and in [14] for the billiard system. In section 4 we draw the major conclusions and present an outlook.

2. Billiard

The billiard domain $\mathcal{B}$ under study is defined, as introduced in [45, 46], by the quadratic conformal map of the unit circle $|z| = 1$ of the z-complex plane onto the boundary $\partial\mathcal{B}$ in the w-complex plane (which is the physical plane) as follows

$$w = z + \lambda z^2. \tag{4}$$

We choose $\lambda = 0.15$, in which case the boundary is convex and the dynamics is of the mixed-type. There are regions of regular motion near the boundary (Lazutkin's caustics) and also in the interior [47, 48, 49]. Classical dynamics is fully determined by the bounce map on the phase space cylinder (s, p), where the arclength parameter $s \in [0, \mathcal{L})$ determines the position of the collision point on the boundary, whilst $p \in [-1, +1]$ is the sine of the reflection angle and is thus the momentum conjugate to s. In fact, due to the reflection symmetry and the time-reversal symmetry, one quadrant $(s, p) \in [0, \mathcal{L}/2] \times [0, 1]$ is enough to be presented. By ρ_1 we denote the relative phase space volume (not to be confused with the area on the Poincaré surface of section) of the classical regular components and by $\rho_2 = 1 - \rho_1$ its complement, the relative volume of the chaotic component. In our case $\rho_1 = 0.175$ [13].

The quantum mechanics is determined by the stationary Schrödinger equation which in appropriate units is just the Helmholtz equation for the billiard domain $\mathcal{B}$,

$$\Delta\psi + k^2\psi = 0, \tag{5}$$

with the Dirichlet boundary condition $\psi = 0$ on the boundary $\partial\mathcal{B}$. The energy eigenvalues $E_j = k_j^2$, $j = 1, 2, 3, \ldots$ define the energy spectrum of the billiard, and after spectral unfolding, based on the Weyl formula with perimeter corrections (see e.g. [1]), we study its statistical properties. The total spectrum can be conceptually decomposed into regular and chaotic eigenstates. The further we are in the semiclassical regime, the "cleaner" is this separation. The physical separation is possible by looking at the structure of the Wigner functions [4], or better, Husimi functions [5] (which are Gaussian averaged Wigner functions). In doing this for a great number of eigenstates we first introduce quantum representation analogous to the classical one. To this end we remind that the *boundary function* $u(s)$, which is the normal derivative of $\psi(\mathbf{r})$ on the boundary at the point s,

$$u(s) = \mathbf{n} \cdot \nabla_{\mathbf{r}} \psi\left(\mathbf{r}(s)\right), \tag{6}$$

where $\mathbf{n}$ is the unit outward vector normal to the boundary at position s, uniquely determines the solution $\psi(\mathbf{r})$ at any point in the interior of $\mathcal{B}$, by the relation (see e.g. [50])

$$\psi_j(\mathbf{r}) = -\oint dt\; u_j(t)\; G\left(\mathbf{r}, \mathbf{r}(t)\right). \tag{7}$$

Thus, in certain analogy to the classical mechanics, the quantum mechanics is completely described by the boundary functions $u_j(s)$. Moreover, the eigenvalue problem (5) is equivalent to the following integral equation

$$u(s) = -2\oint dt\; u(t)\; \mathbf{n} \cdot \nabla_{\mathbf{r}} G(\mathbf{r}, \mathbf{r}(t)). \tag{8}$$

Here $\mathbf{r}(t)$ is the position vector at the point $s = t$ on the boundary, whilst $\mathbf{r}$ is the position vector inside the billiard $\mathcal{B}$. $G(\mathbf{r}, \mathbf{r}(t))$ is the free particle Green function, namely $G(\mathbf{r}, \mathbf{r}') = -\frac{i}{4} H_0^{(1)}(k|\mathbf{r} - \mathbf{r}'|)$, where $H_0^{(1)}(x)$ is the zero order Hankel function of the first kind.

Having set up this representation we proceed by defining the *Poincaré Husimi functions*, following the formalism as in [51], which in turn is based on [52] - [55], namely we define the manifestly s-periodic coherent states (with the period $\mathcal{L}$)

$$c_{(q,p),k}(s) = \sum_{m\in\mathbb{Z}} \exp\{i\, k\, p\, (s - q + m\mathcal{L})\} \exp\left(-\frac{k}{2}(s - q + m\mathcal{L})^2\right). \tag{9}$$

They are concentrated at $(q, p) \in [0, \mathcal{L}] \times \mathbb{R}$. Here we have omitted all normalization factors, because in the end we shall normalize the Poincaré Husimi functions anyway. Then, using this, the Poincaré Husimi function associated with the j-th eigenstate represented by the boundary function $u_j(s)$ with the eigenvalue $k = k_j$, is

$$H_j(q, p) = \left| \int_{\partial\mathcal{B}} c_{(q,p),k_j}(s)\; u_j(s)\; ds \right|^2, \tag{10}$$

which is positive definite by construction. In the semiclassical limit $j \to \infty$, and $k_j \to \infty$, we shall observe that the Poincaré Husimi function is concentrated on the classical invariant regions, which can be an invariant torus, a chaotic component, or the entire Poincaré surface of section (s, p) if the motion is ergodic.



Figure 1. Examples of a chaotic (left) and a regular (right) state in the Poincaré Husimi representation. The corresponding parameter values are: $k_j = 2000.0181794$, $M_j = 0.981$ and $k_j = 2000.0777155$, $M_j = -0.821$. The gray background is the classically chaotic invariant component. We show only one quarter of the surface of section $(s, p) \in [0, \mathcal{L}/2] \times [0, 1]$, because due to the reflection symmetry and time-reversal symmetry the four quadrants are equivalent.

The quantum eigenstates were calculated using the method of Vergini and Saraceno [55] with great accuracy, for all eigenstates (587654) within the interval $k \in [2000, 2500]$. The number of eigenstates below $k = 2000$ is estimated by the Weyl rule as about 1.000.000. Then, when calculating the Poincaré Husimi functions, the momentum p is rescaled by the eigenvalue k_j, such that $p = 1$ corresponds to the original $p = k_j$. Two examples of the Poincaré Husimi functions are shown in figure 1.

Next the Poincaré Husimi function $H(q, p)$ (10) was calculated on the grid points (400×400) on one quadrant $(s, p) \in [0, \mathcal{L}/2]\times[0, 1]$ for each eigenstate j, then normalized such that the sum over all grid points is equal to one, and the overlap index M was calculated according to the definition

$$M = \sum_{i,j} H_{i,j}\, A_{i,j}. \tag{11}$$

Here, $A_{i,j}$ is equal to $+1$ if the grid point (i, j) belongs to the chaotic region, and -1 if it belongs to the regular region. Therefore, due to the normalization of $H_{i,j}$, and in the ideal (semiclassical case) M is either $+1$ or -1. In practice, M is not exactly $+1$ or -1, but can have a value in between. The reasons are two, first the finite discretization of the phase space (the finite size grid), and second, the finite wavelength (not sufficiently small effective Planck constant, for which we can take just $1/k_j$). If so, the question is, where to cut the distribution of the M-values, at the threshold value M_t, such that all states with $M < M_t$ are declared regular and those with $M > M_t$ chaotic.

There are two natural criteria: **(I)** *The classical criterion:* the threshold value M_t is chosen such that we have exactly ρ_1 fraction of regular levels and $\rho_2 = 1 - \rho_1$ of chaotic levels. **(II)** *The quantum criterion:* we choose M_t such that we get the best possible agreement of the chaotic level spacing distribution with the Brody distribution, which is expected to capture the dynamical localization effects of the chaotic eigenstates.

In figure 2 we show the level spacing distribution after separation using the classical criterion. In figure 3 we show the U-function plot (introduced in [9], and explained in detail in [13]), demonstrating that Brody distribution is indeed excellent description of the chaotic level spacings, in case of the criterion (II) even better than using the criterion



Figure 2. Separation of levels using the classical criterion $M_t = 0.431$. (a; left) The level spacing distribution for the chaotic subspectrum after unfolding, in perfect agreement with the Brody distribution $\beta = 0.444$. (b; right) The level spacing distribution for the regular part of the spectrum, after unfolding, in excellent agreement with Poisson.



Figure 3. The U-function plots as differences $U(data) - U(ideal)$ for the regular and chaotic levels, for both criteria, the classical one $M_t = 0.431$ and the quantum one $M_t = 0.75$. The belts around the data lines indicate the expected statistical ± one-sigma errors.

(I). Also, Poisson distribution of the regular levels is clearly manifested. Finally, we show in figure 4 that the Izrailev level spacing distribution [30, 33, 29] (3) is much less significant than the Brody distribution.

3. Kicked rotator

One of the main models of time-periodic (Floquet) systems is the kicked rotator introduced by Casati, Chirikov, Ford and Izrailev in 1979 [28]. For an extensive review



Figure 4. We show the U-function plot for the chaotic levels, clearly showing that Brody distribution (dashed) is much better, even perfect, description of the U-function than the Izrailev. Left is the case when Brody gives the best posssible fit with Izrailev at the same quantum threshold $M_t = 0.75$, whilst right is the case when Izrailev gives the best possible fit with the Brody at the same threshold $M_t = -0.5$. This is impressive, because the effects are small, and the statistical significance very high. The belt around the data line indicates the expected statistical ± one-sigma error.

see [29]. The Hamiltonian is

$$H = \frac{p^2}{2I} + V_0\, \delta_T(t)\, \cos\theta. \tag{12}$$

Here p is the (angular) momentum, I the moment of inertia, V_0 is the strength of the periodic kicking, θ is the (canonically conjugate, rotation) angle, and $\delta_T(t)$ is the periodic Dirac delta function with period T. Since between the kicks the rotation is free, the Hamilton equations of motion can be immediately integrated, and thus the dynamics can be reduced to the standard mapping, or so-called Chirikov-Taylor mapping, given by

$$P_{n+1} = P_n + K \sin\theta_{n+1}, \quad \theta_{n+1} = \theta_n + P_n, \tag{13}$$

where $P_n = p_n T/I$, and the system is now governed by a single classical *dimensionless* control parameter $K = V_0 T/I$, and the mapping is area preserving. The quantities (θ_n, p_n) refer to their values just immediately after the n-th kick.

The quantum kicked rotator (QKR) is the quantized version of (12), namely

$$\hat{H} = -\frac{\hbar^2}{2I}\frac{\partial^2}{\partial\theta^2} + V_0\, \delta_T(t)\, \cos\theta. \tag{14}$$

For such a Floquet system the Floquet operator $\hat{F}$ acting on the wavefunctions $\psi(\theta)$, $\theta \in [0, 2\pi)$, upon each period (of length T) can be written as (see e.g. [1], Sec. 4)

$$\hat{F} = \exp\left(-ik\cos\theta\right)\exp\left(-\frac{i\tau}{2}\frac{\partial^2}{\partial\theta^2}\right), \tag{15}$$

where now we have obviously two *dimensionless* quantum control parameters $k = V_0/\hbar$, $\tau = \hbar\, T/I$, which satisfy the relationship $K = k\tau = V_0T/I$, K being the classical *dimensionless* control parameter of (13). We have investigated in detail the classical and quantal system at $K = 7$, which is classically fully chaotic regime (no islands of stability), in contradistinction to the usually employed value $K = 5$, which still is of the mixed-type (regular regions have relative size/area of order of 2%). Many more cases of $K > 7$ have been studied extensively in [34]. If $\tau/(4\pi)$ is a rational number, then we have the so-called *quantum resonance* and continuous quasienergy spectrum. If $\tau/(4\pi)$ is an irrational number, we face the dynamical localization which sets in when the evolution time exceeds the Heisenberg time t_H. The infinite dimensional system exhibits in such cases the dynamical localization and consequently the Poisson level spacing distribution, for the quasienergies. However, if we study the finite dimensional system, which can be regarded as one of the possible discretizations of the kicked rotator (or of standard map), the scenario changes completely. Then it is convenient to work in the angular momentum basis, defined by $|n\rangle = \exp(in\theta)$. We have studied the following system

$$u_n(t+T) = \sum_{m=1}^{N} U_{nm} u_m(t), \qquad n,m = 1,2,...,N\ . \tag{16}$$

The finite symmetric unitary matrix U_{nm} determines the evolution of an N-dimensional vector, namely the Fourier transform $u_n(t)$ of $\psi(\theta,t)$, and is composed in the following way

$$U_{nm} = \sum_{n'm'} G_{nm'} B_{n'm'} G_{n'm}, \tag{17}$$

where $G_{ll'} = \exp\left(i\tau l^2/4\right)\delta_{ll'}$ is a diagonal matrix corresponding to free rotation during a half period $T/2$, and the matrix $B_{n'm'}$ describing the one kick has the following form

$$B_{n'm'} = \frac{1}{2N+1} \sum_{l=1}^{2N+1} \left\{ \cos\left[(n'-m')\,\frac{2\pi l}{2N+1}\right] - \cos\left[(n'+m')\frac{2\pi l}{2N+1}\right]\right\} \\ \times \exp\left[-ik\cos\left(\frac{2\pi l}{2N+1}\right)\right]. \tag{18}$$

The model (16-18) with a finite number of states is considered as the quantum analogue of the classical standard mapping on the torus with closed momentum p and phase θ, where U_{nm} describes only the odd states of the systems, i.e. $\psi(\theta) = -\psi(-\theta)$, provided we have the case of the quantum resonance, namely $\tau = 4\pi r/(2N+1)$, where r is a positive integer. The matrix (18) is obtained by starting the derivation from the odd-parity basis of $\sin(n\theta)$ rather than the general angular momentum basis $\exp(in\theta)$.

Nevertheless, we use this model for any value of τ and k, as a model which in the resonant and in the generic case (irrational $\tau/(4\pi)$) corresponds to the classical kicked rotator, and in the limit $N \to \infty$ approaches the infinite dimensional model, restricted to the symmetry class of the odd eigenfunctions. It is of course just one of the possible discrete approximations to the continuous infinite dimensional model.



Figure 5. Intermediate statistics (panel (a)) for distribution $P(S)$ (histogram - black solid line) of the model [Eqs. (16)-(17)] fitted with distribution $P_B(S)$ (black dashed line) and $P_I(S)$ (black dotted line) for 641 matrices of size 398, with $K = 7$ and k in small interval around 11. The gray lines indicate the two extreme distributions, i.e. the Poisson and Wigner. In panels (b)and (c) we show the difference of the numerical data and the best fitting Brody (black line) and Izrailev (gray line) PDFs by using the U-function and W-distribution. Thus in case of the ideal fitting the data would lie on the abscissa. In this case, based on the $P(S)$ fit we get $\beta = 0.421$ for Brody and $\beta = 0.416$ for Izrailev fit, whilst based on the $W(S)$ fit we get $\beta = 0.421$ and $\beta = 0.401$, correspondingly.

In figure 5 we show the main results. We clearly see that the Brody distribution is an excellent fit to the level spacing distribution, even more clearly in the U-function plots, and we also see that it is empirically better than Izrailev's fit. All other details and plentiful results will be published in a separate paper [34], where we find that Brody distribution is better than Izrailev for almost all values of β, at $K = 7$.

4. Conclusions

We have demonstrated that in case of dynamical localization of chaotic eigenstates the corresponding energy spectrum obeys very accurately the Brody distribution, so we observe the fractional power law level repulsion with the exponent $\beta \in [0, 1]$, which is the Brody parameter. This has been demonstrated for the finite dimensional kicked rotator (quantized standard map) and its quasienergy spectra, as well as for the chaotic eigenstates in a time-independent Hamilton system, namely the convex billiard of the mixed-type (Robnik 1983, $\lambda = 0.15$; [45, 46]), and the corresponding energy levels. The separation of the regular and chaotic eigenstates has been performed by means of Poincaré Husimi functions of the boundary function, projected onto the classical Poincaré surface of section. The number of objects in the histogram is 587653, and thus the statistical significance is great. The successful separation of course also confirms the Berry-Robnik picture [7] of separating the regular and chaotic levels in the semiclassical limit, where the tunneling effects can be neglected. Many more details and further results will be published in two separate papers [34] and [14].

The theoretical derivation of the Brody level spacing distribution for the dynamically localized chaotic eigenstates is an open problem for the future. The billiard systems are not just nice theoretical toy models, but are suitable also for the experimental applications, like e.g. in quantum dots, and microwave cavities introduced and studied extensively over decades by H.-J. Stöckmann [1]. We also propose to study from the present point of view the hydrogen atom in strong magnetic field as an example of classical and quantum chaos par excellence, as introduced in [56, 57, 58, 59, 60], although the technical efforts to obtain large stretches of high-lying eigenstates and the corresponding energy levels are much bigger than in billiard systems, where we have a great number of different elegant numerical techniques [61], all of them used in our recent work [13].

Acknowledgements

Financial support of the Slovenian Research Agency ARRS under the grants P1-0306 and J1-4004 is gratefully acknowledged. T.M. was also supported by the European Union (European Social Fund) and Greek national funds through the Operational Program “Education and Lifelong Learning” of the National Strategic Reference Framework (NSRF) - Research Funding Program THALES: Investing in knowledge society through the European Social Fund.

References

[1] Stöckmann H.-J. 1999 *Quantum Chaos - An Introduction* (Cambridge: Cambridge University Press).
[2] Haake F 2001 *Quantum Signatures of Chaos* (Berlin: Springer)
[3] Robnik M 1998 *Nonl. Phen. in Compl. Syst.* (Minsk) **1** 1
[4] Wigner E P 1932 *Phys. Rev.* **40** 749
[5] Husimi K 1940 *Proc. Phys. Math.. Soc. Jpn.* **22** 264
[6] Percival I C 1973 *J. Phys. B: At. Mol. Phys.* **6** L229
[7] Berry M V and Robnik M 1984 *J. Phys. A: Math. Gen.* **17** 2413
[8] Prosen T and Robnik M 1994 *J. Phys. A: Math. Gen.* **27** L459
[9] Prosen T and Robnik M 1994 *J. Phys. A: Math. Gen.* **27** 8059
[10] Prosen T and Robnik M 1999 *J. Phys. A: Math. Gen.* **32** 1863
[11] Prosen T 1998 *J. Phys. A: Math. Gen.* **31** L345
[12] Prosen T 1998 *J. Phys. A: Math. Gen.* **31** 7023

[13] Batistić B and Robnik M 2010 *J. Phys. A: Math. Theor.* **43** 215101
[14] Batistić B and Robnik M 2013 *J. Phys. A: Math. Theor.* to be submitted
[15] Mehta M L 1991 *Random Matrices* (Boston: Academic Press)
[16] Guhr T, Müller-Groeling A and Weidenmüller H A 1998 *Phys. Rep.* **299** Nos. 4-6 189
[17] Bohigas O, Giannoni M.-J and Schmit C 1984 *Phys. Rev. Lett.* **52** 1
[18] Casati G, Valz-Gris F and Guarneri I 1980 *Lett. Nuovo Cimento* **28** 279
[19] Robnik M and Berry M V 1986 *J. Phys. A: Math. Gen.* **19** 669
[20] Robnik M 1986 *Lect. Notes Phys.* **263** 120
[21] Berry M V 1985 *Proc. Roy. Soc. Lond. A* **400** 229
[22] Gutzwiller M C 1990 *Chaos in Classical and Quantum Mechanics* (Berlin: Springer) and references therein
[23] Sieber M and Richter K 2001 *Phys. Scr.* **T90** 128
[24] Müller S, Heusler S, Braun P, Haake F and Altland A 2004 *Phys. Rev. Lett.* **93** 014103
[25] Heusler S, Müller S, Braun P and Haake F 2004 *J. Phys.A: Math. Gen.* **37** L31
[26] Müller S, Heusler S, Braun P, Haake F and Altland A 2005 *Phys. Rev. E* **72** 046207
[27] Müller S, Heusler S, Altland A, Braun P, and Haake F 2009 *New J. of Phys.* **11** 103025
[28] Casati G, Chirikov B, Ford J and Izrailev F M 1979 *Lect. Notes Phys.* **93** 334
[29] Izrailev F M 1990 *Phys. Rep.* **196** 299
[30] Izrailev F M 1988 *Phys. Lett. A* **134** 13
[31] Izrailev F M 1986 *Phys. Rev. Lett.* **56** 541
[32] Izrailev F M 1987*Phys. Lett. A* **125** 250
[33] Izrailev F M 1989 *J. Phys. A: Math. Gen.* **22** 865
[34] Manos T and Robnik M 2013 *submitted* (preprint ArXiv: 1301.4187)
[35] Prosen T 2000 in *Proceedings of the International School of Physics "Enrico Fermi", Course CXLIII, Eds. G. Casati, I. Guarneri and U. Smilyanski* (Amsterdam: IOS Press) p 473
[36] Li Baowen and Robnik M 1995 *J. Phys. A: Math. Gen.* **28** 4843
[37] Robnik M and Veble G 1998 *J. Phys. A: Math. Theor.* **31** 4669
[38] Berry M V 1977 *J Phys. A: Math. Gen.* **12** 2083
[39] Veble G, Robnik M and Liu Junxian 1999 *J. Phys. A: Math. Gen.* **32** 6423
[40] Brody T A 1973 *Lett. Nuovo Cimento* **7** 482
[41] Brody T A, Flores J, French J B, Mello P A, Pandey A and Wong S S M 1981 *Rev. Mod. Phys.* **53** 385
[42] Vidmar G, Stöckmann H.-J., Robnik M, Kuhl U, Höhmann R and Grossmann S 2007 *J. Phys. A: Math. Theor.* **40** 13883
[43] Grossmann S and Robnik M 2007 *J. Phys. A: Math. Theor.* **40** 409
[44] Grossmann S and Robnik M 2007 *Z. Naturforschung A* **62** 471
[45] Robnik M 1983 *J. Phys. A: Math. Gen.* **16** 3971
[46] Robnik M 1984 *J. Phys. A: Math. Gen.* **17** 1049
[47] Lazutkin V F 1981 *Convex billiard and eigenfunctions of the Laplace operator* (University of Leningrad) in Russian
[48] Lazutkin V F 1991 *KAM Theory and Semiclassical Approximations to Eigenfunctions* (Berlin: Springer)
[49] Hayli A, Dumont T, Moulin-Ollagier J and Strelcyn J M 1987 *J. Phys. A: Math. Gen.* **20** 3237
[50] Berry M V and Wilkinson M 1984 *Proc. Roy. Soc. Lond. A* **392** 15
[51] Bäcker A, Fürstberger S and Schubert R 2004 *Phys. Rev. E* **70** 036204
[52] Crespi B, Perez G and Chang S.-J. 1993 *Phys. Rev. E* **47** 986
[53] Tualle J M and Voros A 1995 *Chaos, Solitons and Fractals* **5** 1085
[54] Simonotti F P, Vergini E and Saraceno M 1997 *Phys. Rev. E* **56** 3859
[55] Vergini E and Saraceno M 1995 *Phys. Rev. E* **52** 2204
[56] Robnik M 1981 *J. Phys. A: Math. Gen.* **14** 3195-3216
[57] Robnik M 1982 *J. Phys. Colloque C2* **43** 29

[58] Hasegawa H, Robnik M and Wunner G 1989 *Prog. Theor. Phys. Suppl. (Kyoto)* **98** 198-286

[59] Wintgen D and Friedrich H 1989 *Phys. Rep.* **183** 38

[60] Ruder H, Wunner G, Herold H and Geyer F 1994 *Atoms in Strong Magnetic Fields* (Heidelberg: Springer)

[61] Veble G, Prosen T and Robnik M 2007 *New J. Phys.* **9** 1